




The Most Prescribed IV Iron in the US¹

Venofer[®]

iron sucrose injection, USP


*The Most Prescribed
IV Iron in the US¹*

Venofer is indicated for the treatment of iron deficiency anemia (IDA) in patients with chronic kidney disease (CKD).

Venofer must only be administered intravenously, either by slow injection or infusion.

**Dosing and administration
guide**

Guidance for management of anemia in non-dialysis-dependent chronic kidney disease (NDD-CKD)*



See important Notice on page S12 of the Clinical Practice Guideline regarding appropriate usage and limitations. Health care providers are responsible for exercising their own clinical judgment when considering the recommendations set forth in the guidelines.

*Adapted from the KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. Official Journal of the International Society of Nephrology. *Kidney Int.* (2026) 109 (Suppl 1S), S1–S99.

†See page 4 for information about Venofer dosing

7-day stability after dilution

Venofer® (iron sucrose) injection, USP offers 7-day stability in both plastic syringes and IV admixtures diluted with 0.9% sodium chloride (NaCl) injection, helping to avoid potential waste.

	Concentration	Controlled room temperature (25°C ± 2°C)	Refrigerated (4°C ± 2°C)
Plastic syringe	2 mg to 10 mg elemental iron per mL 20 mg of elemental iron per mL (undiluted)		
IV admixture (PVC and non-PVC)	1 mg to 2 mg elemental iron per mL		

- Do not dilute in concentrations below 1 mg/mL
- Do not mix Venofer with other medications
- Do not add Venofer to parenteral nutrition solutions
- Do not administer to patients with iron overload

For Intravenous Use Only

INDICATION AND USAGE

Venofer® (iron sucrose) injection, USP is indicated for the treatment of iron deficiency anemia (IDA) in patients with chronic kidney disease (CKD).

IMPORTANT SAFETY INFORMATION

DOSAGE AND ADMINISTRATION

Pediatric Patients (2 Years of Age and Older)

The dosing for iron replacement treatment in pediatric patients with Peritoneal or Hemodialysis-Dependent CKD or Non-Dialysis Dependent CKD have not been established.

CONTRAINDICATIONS

Known hypersensitivity to Venofer.

Please see additional Important Safety Information throughout, and accompanying [Full Prescribing Information](#).

Dosing and administration for adult CKD patients²

The usual adult total treatment course of Venofer® (iron sucrose) injection, USP is 1,000 mg. Venofer treatment may be repeated if iron deficiency reoccurs.

IV push	100 mg over 2 to 5 minutes	• Hemodialysis-dependent chronic kidney disease (HDD-CKD)* • Consecutive dialysis sessions
	200 mg over 2 to 5 minutes	• Non-dialysis-dependent chronic kidney disease (NDD-CKD) • Five occasions over 14 days
Infusion	100 mg in a maximum of 100 mL over at least 15 minutes	• Hemodialysis-dependent chronic kidney disease (HDD-CKD)* • Consecutive dialysis sessions
	200 mg in a maximum of 100 mL over 15 minutes	• Non-dialysis-dependent chronic kidney disease (NDD-CKD)* • Five occasions over 14 days
	2 infusions each of 300 mg in a maximum of 250 mL over 1.5 hrs followed by 1 dose of 400 mg in a maximum of 250 mL over 2.5 hrs	• Peritoneal-dialysis-dependent chronic kidney disease (PDD-CKD) • 3 divided infusions each within a 28-day period • 14 days apart

*Administer early during the dialysis session.

IMPORTANT SAFETY INFORMATION (Cont'd)

WARNINGS AND PRECAUTIONS

Hypersensitivity Reactions: Serious hypersensitivity reactions, including anaphylactic-type reactions, some of which have been life-threatening and fatal, have been reported in patients receiving Venofer. Patients may present with shock, clinically significant hypotension, loss of consciousness, and/or collapse. If hypersensitivity reactions or signs of intolerance occur during administration, stop Venofer immediately. Monitor patients for signs and symptoms of hypersensitivity during and after Venofer administration for at least 30 minutes and until clinically stable following completion of the infusion. Only administer Venofer when personnel and therapies are immediately available for the treatment of serious hypersensitivity reactions. Most reactions associated with intravenous iron preparations occur within 30 minutes of the completion of the infusion.

Hypotension: Venofer may cause clinically significant hypotension. Monitor for signs and symptoms of hypotension following each administration of Venofer. Hypotension following administration of Venofer may be related to rate of administration and/or total dose delivered.

Dosing and administration for pediatric CKD patients²

Pediatric patients

(2 years of age or older) with HDD-CKD

For iron maintenance treatment, administer Venofer® (iron sucrose) injection, USP

- At a dose of 0.5 mg/kg, not to exceed 100 mg per dose
- Every 2 weeks for 12 weeks
- Undiluted by slow intravenous injection over 5 minutes or diluted in 0.9% NaCl at concentrations of 1 to 2 mg/mL and administered over 5 to 60 minutes

Do not dilute to concentrations below 1 mg/mL

Pediatric patients

(2 years of age or older) with NDD-CKD or PDD-CKD who are on erythropoietin therapy for iron maintenance treatment

- For iron maintenance treatment, administer Venofer
- At a dose of 0.5 mg/kg, not to exceed 100 mg per dose
- Every 4 weeks for 12 weeks
- Undiluted by slow intravenous injection over 5 minutes or diluted in 0.9% NaCl at concentrations of 1 to 2 mg/mL and administered over 5 to 60 minutes

Do not dilute to concentrations below 1 mg/mL

Venofer treatment may be repeated if necessary. The dosing for iron replacement treatment in pediatric patients with HDD-CKD, NDD-CKD, or PDD-CKD has not been established.

Venofer has not been studied in patients younger than 2 years old.

IMPORTANT SAFETY INFORMATION (Cont'd)

WARNINGS AND PRECAUTIONS

Iron Overload: Excessive therapy with parenteral iron can lead to excess storage of iron with the possibility of iatrogenic hemosiderosis. All adult and pediatric patients receiving Venofer require periodic monitoring of hematologic and iron parameters (hemoglobin, hematocrit, serum ferritin and transferrin saturation). Do not administer Venofer to patients with evidence of iron overload. Transferrin saturation (TSAT) values increase rapidly after intravenous administration of iron sucrose; do not perform serum iron measurements for at least 48 hours after intravenous dosing.

Please see additional Important Safety Information throughout, and accompanying [Full Prescribing Information](#).

For Intravenous Use Only

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

Known hypersensitivity to Venofer.

WARNINGS AND PRECAUTIONS

Hypersensitivity Reactions: Serious hypersensitivity reactions, including anaphylactic-type reactions, some of which have been life-threatening and fatal, have been reported in patients receiving Venofer. Patients may present with shock, clinically significant hypotension, loss of consciousness, and/or collapse. If hypersensitivity reactions or signs of intolerance occur during administration, stop Venofer immediately. Monitor patients for signs and symptoms of hypersensitivity during and after Venofer administration for at least 30 minutes and until clinically stable following completion of the infusion. Only administer Venofer when personnel and therapies are immediately available for the treatment of serious hypersensitivity reactions. Most reactions associated with intravenous iron preparations occur within 30 minutes of the completion of the infusion.

Hypotension: Venofer may cause clinically significant hypotension. Monitor for signs and symptoms of hypotension following each administration of Venofer. Hypotension following administration of Venofer may be related to rate of administration and/or total dose delivered.

Iron Overload: Excessive therapy with parenteral iron can lead to excess storage of iron with the possibility of iatrogenic hemosiderosis. All adult and pediatric patients receiving Venofer require periodic monitoring of hematologic and iron parameters (hemoglobin, hematocrit, serum ferritin, and transferrin saturation). Do not administer Venofer to patients with evidence of iron overload. Transferrin saturation (TSAT) values increase rapidly after intravenous administration of iron sucrose; do not perform serum iron measurements for at least 48 hours after intravenous dosing.

ADVERSE REACTIONS

Adult Patients: The most common adverse reactions in clinical trials ($\geq 2\%$ and greater than comparator) included diarrhea, nausea, vomiting, headache, dizziness, hypotension, pruritus, pain in extremity, arthralgia, back pain, muscle cramp, injection site reactions, chest pain, and peripheral edema.

Pediatric Patients: The most common adverse reactions in clinical trials ($\geq 2\%$) were headache, respiratory tract viral infection, peritonitis, vomiting, pyrexia, dizziness, cough, nausea, arteriovenous fistula thrombosis, hypotension, and hypertension.

Post-Marketing Experience

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. In post-marketing safety studies of Venofer in 1,051 patients with hemodialysis-dependent chronic kidney disease (HDD-CKD), adverse reactions reported by $> 1\%$ were cardiac failure congestive, sepsis, and dysgeusia.

- *Immune system disorders:* anaphylactic-type reactions, angioedema
- *Psychiatric disorders:* confusion
- *Nervous system disorders:* convulsions, collapse, light-headedness, loss-of-consciousness

- *Cardiovascular system*: bradycardia, shock, acute myocardial ischemia with or without myocardial infarction or with in-stent thrombosis in the context of a hypersensitivity reaction
- *Respiratory, thoracic, and mediastinal disorders*: bronchospasm, dyspnea
- *Musculoskeletal and connective tissue disorders*: back pain, swelling of the joints
- *Renal and urinary disorders*: chromaturia
- *General disorders and administration site conditions*: hyperhidrosis

Symptoms associated with Venofer total dosage or infusing too rapidly included hypotension, dyspnea, headache, vomiting, nausea, dizziness, joint aches, paresthesia, abdominal and muscle pain, edema, and cardiovascular collapse. These adverse reactions have occurred up to 30 minutes after the administration of Venofer injection. Reactions have occurred following the first dose or subsequent doses of Venofer. Slowing the infusion rate may alleviate symptoms.

Injection site discoloration has been reported following extravasation. Assure stable intravenous access to avoid extravasation.

DRUG INTERACTIONS

Venofer may reduce the absorption of concomitantly administered oral iron preparations.

USE IN SPECIFIC POPULATIONS

Pregnancy

Untreated IDA in pregnancy is associated with adverse maternal outcomes such as post-partum anemia. Adverse pregnancy outcomes associated with IDA include increased risk for preterm delivery and low birth weight.

Severe adverse reactions including circulatory failure (severe hypotension, shock including in the context of anaphylactic reaction) may occur in pregnant women with parenteral iron products (such as Venofer), which may cause fetal bradycardia, especially during the second and third trimester.

Pediatric Use

Safety and effectiveness of Venofer for iron replacement treatment in pediatric patients with dialysis-dependent or non-dialysis-dependent CKD have not been established.

Geriatric Use

Dose administration to an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

INDICATION AND USAGE

Venofer® (iron sucrose) injection, USP is indicated for the treatment of iron deficiency anemia (IDA) in patients with chronic kidney disease (CKD).

For additional Safety Information, please see [Full Prescribing Information](#).

You are encouraged to report adverse drug events to American Regent, Inc. at 1-800-734-9236 or to the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.

The Most Prescribed Iron in the US¹

Available in the following sizes

Pack NDC	Strength (Each mL contains 20 mg of elemental iron)	Supplied as
0517-2310-05	200 mg	10 mL Single-dose vial
0517-2325-10	50 mg	2.5 mL Single-dose vial
0517-2340-10	100 mg	5 mL Single-dose vial
0517-2340-25	100 mg	5 mL Single-dose vial

Millions prescribed. Millions treated.*



**You are encouraged to report adverse drug events (ADEs)
to American Regent:**

T 1.800.734.9236; **E** pv@americanregent.com; **F** 1.610.650.0170

ADEs may also be reported to the FDA:
1.800.FDA.1088 or www.fda.gov/medwatch

Medical information:
T 1.888.354.4855
(9:00 am–5:00 pm Eastern Time, Monday–Friday)
www.americanregent.com/medical-affairs

1-800-645-1706 \ \ AMERICANREAGENT.COM



REFERENCES:

1. IQVIA. NSP Audit from January 2020 to January 2025.
2. Venofer[®] (iron sucrose) injection, USP. Package insert. American Regent, Inc.

Venofer[®] and the Venofer logo are registered trademarks of Vifor (International), Inc., Switzerland. Millions prescribed. Millions treated.[®] is a registered trademark of American Regent, Inc. Venofer[®] is manufactured under license from Vifor (International) Inc., Switzerland.